

Search for the QCD Critical Point by Higher Moments of Net-proton Multiplicity Distributions at STAR

Xiaofeng Luo (for the STAR Collaboration)

*Key Laboratory of Quark&Lepton (MOE) and Institute of Particle Physics,
Central China Normal University, 430079 Wuhan, China*

Abstract

We present the measurements of the higher moments of the net-proton multiplicity distributions for Au+Au collisions from the first phase of the beam energy scan program at RHIC. The measurements are carried out at midrapidity ($|y| < 0.5$) and within the transverse momentum range $0.4 < p_T < 0.8$ GeV/c by STAR. The moment products, $\kappa\sigma^2$ and $S\sigma$, which are related to the ratios of various order baryon number susceptibilities and sensitive to the correlation length, are applied to search for the QCD critical point. We report the centrality and energy dependence of the moment products ($\kappa\sigma^2$ and $S\sigma$), results are compared with Poisson expectations. The experiment data are also compared with transport model calculation to understand the non-critical physics effects.

1. Introduction

In the year 2010, the Relativistic Heavy Ion Collider (RHIC) launched the Beam Energy Scan (BES) program [1] and tuned the Au+Au collision energy from $\sqrt{s_{NN}}=200$ GeV down to 7.7 GeV. The main goal of the BES program is to look for the signature of the first order phase transition and Quantum Chromodynamics (QCD) Critical Point (CP) [2, 3, 4, 5]. By varying the colliding energy, we can access a broad region of the QCD phase diagram with a baryon chemical potential (μ_B) range from about 20 MeV to 450 MeV [1]. As the diverges of the correlation length (ξ) near the CP, non-monotonic variations of observables with collision energy are believed to be the best signature of CP. Higher moments (skewness (S) and kurtosis (κ) *etc.*) of conserved quantities, such as net-baryon, net-charge and net-strangeness, distributions are more sensitive to the correlation length [6, 7, 8] than the variance (σ^2) and could be detectable if the signature survive in the evolution of the system. On the other hand, the moment products, $\kappa\sigma^2$ and $S\sigma$, are also related to the ratios of various order susceptibilities as in Lattice QCD [9, 10, 11], HRG model [12] and PQM model [13]. Ratios of baryon number susceptibilities used to compare with the experimental data as $\kappa\sigma^2 = \chi_B^{(4)}/\chi_B^{(2)}$ and $S\sigma = \chi_B^{(3)}/\chi_B^{(2)}$ cancel out the volume effect. Theoretical calculations demonstrate that the experimentally measurable net-proton (proton number minus anti-proton number) number fluctuations can effectively reflect the fluctuations of the net-baryon number [14]. In this paper, we will present measurements for higher moments of event-by-event net-proton multiplicity distributions from the first phase of the BES program at RHIC.

2. Results and Discussion

The data of Au+Au collisions at $\sqrt{s_{NN}}=7.7, 11.5, 19.6, 27, 39, 62.4$ and 200 GeV from the first phase of the BES program are used in the analysis. We have also studied $p + p$ collisions at $\sqrt{s_{NN}}=62.4$ and 200 GeV. The protons and anti-protons are identified with the ionization energy loss (dE/dx) measured by Time Projection Chamber (TPC) at midrapidity ($|\eta| < 0.5$) and within the transverse momentum range $0.4 < p_T < 0.8$ GeV/c. To suppress autocorrelation effects between measured net-proton fluctuations and centrality in our results, centralities are determined by the uncorrected charge particle multiplicity ($dN_{ch}/d\eta$) by excluding the protons and anti-protons within pseudorapidity $|\eta| < 0.5$, a new technique add to the moment analysis after QM2011 conference. If protons and anti-protons are independent Poissonian distributions, the net-proton multiplicity will follow the skellam distribution, which is expressed as:

$$P(N) = \left(\frac{M_p}{M_{\bar{p}}}\right)^{N/2} I_N(2\sqrt{M_p M_{\bar{p}}}) \exp[-(M_p + M_{\bar{p}})],$$

where $I_N(x)$ is a modified Bessel function, M_p and $M_{\bar{p}}$ are the measured mean values of protons and anti-protons. The various order cumulants (C_n) are closely connected with the moments, e.g., $C_1 = \langle N \rangle = M$, $C_2 = \langle (\Delta N)^2 \rangle = \sigma^2$, $C_3 = \langle (\Delta N)^3 \rangle = S\sigma^3$, $C_4 = \langle (\Delta N)^4 \rangle - 3\langle (\Delta N)^2 \rangle^2 = \kappa\sigma^4$. If the net-proton follows the skellam distribution, then we have, $S\sigma = C_3/C_2 = (M_p - M_{\bar{p}})/(M_p + M_{\bar{p}})$ and $\kappa\sigma^2 = C_4/C_2 = 1$, which provide the Poisson expectations for the moment products.




Figure 1: (Color Online) Centrality dependence of the cumulants of net-proton distributions for Au+Au collisions at $\sqrt{s_{NN}}=7.7, 11.5, 19.6, 27, 39, 62.4$ and 200 GeV. The lines are linear fit to these cumulants. Errors bars are statistical and caps are systematic errors.




Figure 2: (Color Online) Centrality dependence of $S\sigma/\text{Poisson}$ and $\kappa\sigma^2$ of net-proton distributions for Au+Au collisions at $\sqrt{s_{NN}}=7.7, 11.5, 19.6, 27, 39, 62.4$ and 200 GeV. The error bars are statistical and caps are systematic errors.

The various order cumulants ($C_1 - C_4$) shown in Fig. 1 are obtained from the net-proton distributions and correct for the finite centrality bin width effect [4]. The statistical error estimation in Fig.1 are based on the Delta theorem method [15], which is an updated method compare to the analytical method used in QM2011 results [5]. The systematical errors are estimated by varying the track quality cut condition and particle identification. $C_1 - C_4$ values in Fig.1 show

linear increase with N_{part} in each centrality. The lines shown in the Fig.1 are linear fit to those cumulants, which is motivated by the Central Limit Theorem (CLT). In Fig. 1, it is observed that the odd (C_1 and C_3) and even (C_2 and C_4) order cumulants group together. The difference between the two groups are smaller at lower energies. The ratios of the cumulants, which are connected to the moment products as $S\sigma = C_3/C_2$ and $\kappa\sigma^2 = C_4/C_2$, are shown in Fig. 2. It is observed that the $\kappa\sigma^2$ and the $S\sigma$ values normalized to Poisson expectations are below unity for $\sqrt{s_{NN}}$ above 11.5 GeV and above unity for 7.7 GeV in Au+Au collisions. The data presented here also provide information to extract freeze-out conditions in heavy-ion collisions using QCD based approaches [16].




Figure 3: (Color online) Energy dependence of $\kappa\sigma^2$ and $S\sigma$ for net-proton distributions for four collision centralities (0-5%, 5-10%, 30-40% and 70-80%) measured at STAR. The results are compared to UrQMD model calculations and $p + p$ collisions at $\sqrt{s_{NN}}=62.4$ and 200 GeV. The lines in top panel are the Poisson expectations and in the bottom panel shows the $S\sigma$ normalized to the corresponding Poisson expectations.

Fig.3 shows the energy dependence of $\kappa\sigma^2$ and $S\sigma$ of net-proton distributions for four centralities (0-5%, 5-10%, 30-40% and 70-80%) in Au+Au collisions. Bottom panel of Fig.3 shows $S\sigma$ values normalized to the corresponding Poisson expectations. The $\kappa\sigma^2$ and normalized $S\sigma$ values are similar for all of the centralities in Au+Au collisions at $\sqrt{s_{NN}}=39, 62.4$ and 200 GeV and differences are observed between the 0-5% central Au+Au collisions and the peripheral collisions below $\sqrt{s_{NN}}=39$ GeV. The peripheral collision results are closer to unity and for $\sqrt{s_{NN}}=7.7$ GeV, those are above unity. The UrQMD model [17] results are also shown in the Fig. 3 for 0-5% centrality to understand the non-CP effects, such as baryon number conservation and hadronic scattering. The UrQMD calculations show a monotonic decrease with decreasing beam energy.

3. Summary

We present the beam energy ($\sqrt{s_{NN}}=7.7-200$ GeV) and centrality dependence for the higher moments of net-proton distributions in Au+Au collisions from the first phase of the BES program at RHIC. We have discussed the application of those observables to search for the QCD critical point. It is observed that the moment products, $\kappa\sigma^2$ and $S\sigma$, are below the Poisson expectation for central Au+Au collisions. The peripheral collision results are closer to unity and for $\sqrt{s_{NN}}=7.7$ GeV, those are above unity. The experimental data are also compared with the UrQMD model calculation to understand the non-critical physics effects. The UrQMD calculations show a monotonic decrease with decreasing beam energy. Due to the small correlation length and dynamical evolution of the system in the heavy ion collisions, the signature of the CP could be small. This needs comparison between our data and QCD calculations with dynamical evolution of the system. We also need more statistics to get precise measurements below 19.6 GeV and additional data at $\sqrt{s_{NN}}=15$ GeV. These are planned for the second phase of the BES program at RHIC.

Acknowledgement

The work was supported in part by the National Natural Science Foundation of China under grant No. 11205067 and 11135011. CCNU-QLPL Innovation Fund(QLPL2011P01) and China Postdoctoral Science Foundation (2012M511237).

References

References

- [1] M. M. Aggarwal *et al.* (STAR Collaboration), arXiv: 1007.2613.
- [2] M. M. Aggarwal *et al.* (STAR Collaboration), Phys. Rev. Lett. **105**, 022302 (2010).
- [3] X. Luo (for the STAR Collaboration), Acta Phys. Pol. B Proc. Supp. **5**, 497 (2012). [arXiv:1111.5671].
- [4] X. Luo (for the STAR Collaboration), J. Phys.: Conf. Ser. **316**, 012003 (2011).
- [5] B. Mohanty (for the STAR Collaboration), J. Phys. G: Nucl. Part. Phys. **38**, 124023 (2011).
- [6] M. A. Stephanov, Phys. Rev. Lett. **102**, 032301 (2009).
- [7] C. Athanasiou *et al.*, Phys. Rev. D **82**, 074008 (2010).
- [8] M. A. Stephanov, Phys. Rev. Lett. **107**, 052301 (2011).
- [9] R. V. Gavai and S. Gupta, Phys. Lett. B **696**, 459 (2011).
- [10] M. Cheng *et al.*, Phys. Rev. D **79**, 074505 (2009). A. Bazavov, *et al.*, arXiv:1203.0784.
- [11] S. Gupta, X. Luo, B. Mohanty, H. G. Ritter, N. Xu, Science **332**, 1525 (2011).
- [12] F. Karsch and K. Redlich, Phys. Lett. B **695**, 136 (2011).
- [13] V. Skokov *et al.*, J. Phys. G: Nucl. Part. Phys. **38**, 124102 (2011).
- [14] Y. Hatta and M. A. Stephanov, Phys. Rev. Lett. **91**, 102003 (2003).
- [15] X. Luo, J. Phys. G: Nucl. Part. Phys. **39**, 025008 (2012).
- [16] F. Karsch, CPD2011 proceedings, arXiv:1202.4173; A. Bazavov, *et al.*, arXiv:1208.1220.
- [17] M. Bleicher *et al.*, J. Phys. G: Nucl. Part. Phys. **25**, 1859 (1999).